

Leicester

Unitary Authority



This profile was produced on 8 July 2014

Health Profile 2014

Health in summary

The health of people in Leicester is varied compared with the England average. Deprivation is higher than average and about 30.0% (21,000) children live in poverty. Life expectancy for both men and women is lower than the England average.

Living Ionger

Life expectancy is 6.8 years lower for men and 4.5 years lower for women in the most deprived areas of Leicester than in the least deprived areas.

Child health

In Year 6, 21.1% (713) of children are classified as obese, worse than the average for England. The rate of alcohol-specific hospital stays among those under 18 was 25.7*, better than the average for England. This represents 20 stays per year. Levels of teenage pregnancy, GCSE attainment and smoking at time of delivery are worse than the England average.

Adult health

In 2012, 19.6% of adults are classified as obese. The rate of alcohol related harm hospital stays was 717*, worse than the average for England. This represents 1,988 stays per year. The rate of self-harm hospital stays was 137.4*, better than the average for England. This represents 488 stays per year. The rate of smoking related deaths was 293*. This represents 366 deaths per year. Estimated levels of adult physical activity are worse than the England average. Estimated levels of adult excess weight are better than the England average. Rates of sexually transmitted infections and TB are worse than average. The rate of people killed and seriously injured on roads is better than average.

Local priorities

Priorities include young people, premature deaths, supporting independence, mental health and the wider determinants of health. For more information see www.leicester.gov.uk

© Crown Copyright and database rights 2014, Ordnance Survey 100016969 OpenStreetMap contributors ODbL

Population: 332,000

Mid-2012 population estimate. Source: Office for National Statistics.

This profile gives a picture of people's health in Leicester. It is designed to help local government and health services understand their community's needs, so that they can work to improve people's health and reduce health inequalities.

Visit www.healthprofiles.info

or scan this Quick Response code: for more profiles, more information and interactive maps and tools.



Follow @healthprofiles on Twitter


* rate per 100,000 population

Leicester! 2 miles


w.leicester.gov.uk Follow <u>@nealthprofiles</u> on Twitt

Deprivation: a national view

The map shows differences in deprivation levels in this area based on national quintiles (fifths) of the Index of Multiple Deprivation 2010 by Lower Super Output Area. The darkest coloured areas are some of the most deprived areas in England.




This chart shows the percentage of the population in England and this area who live in each of these quintiles.




Life Expectancy: inequalities in this local authority

The charts below show life expectancy for men and women in this local authority for 2010-2012. Each chart is divided into deciles (tenths) by deprivation, from the most deprived decile on the left of the chart to the least deprived decile on the right. The steepness of the slope represents the inequality in life expectancy that is related to deprivation in this local area. If there were no inequality in life expectancy as a result of deprivation, the line would be horizontal.

Life Expectancy Gap for Men: 6.8 years




Life Expectancy Gap for Women: 4.5 years




Health inequalities: changes over time

These charts provide a comparison of the changes in early death rates (in people under 75) between this area and all of England. Early deaths from all causes also show the differences between the most and least deprived quintile in this area. (Data points are the midpoints of 3 year averages of annual rates, for example 2005 represents the period 2004 to 2006).



Health inequalities: ethnicity

Percentage of hospital admissions that were emergencies, by ethnic group



Health Summary for Leicester

The chart below shows how the health of people in this area compares with the rest of England. This area's result for each indicator is shown as a circle. The average rate for England is shown by the black line, which is always at the centre of the chart. The range of results for all local areas in England is shown as a grey bar. A red circle means that this area is significantly worse than England for that indicator; however, a green circle may still indicate an important public health problem.

	icantly worse than England average			England	Regional	average^ England Avera	~9~	Engla
Not significantly different from England average				Worst		25th	75th	
_	icantly better than England average	Local No Per Year	Local	Eng	Eng	Percentile England P.	Percentile	Eng best
Domain	indicator	Per rear	value	value	worst	England R	ange	Desi
Our communities	1 Deprivation	135,783	40.9	20.4	83.8			0.0
	2 Children in poverty (under 16s)	20,955	30.0	20.6	43.6			6.4
	3 Statutory homelessness	69	0.5	2.4	33.2			0.0
	4 GCSE achieved (5A*-C inc. Eng & Maths)	1,905	54.8	60.8	38.1			81.9
	5 Violent crime (violence offences)	5,222	15.8	10.6	27.1			3.3
	6 Long term unemployment	3,753	16.8	9.9	32.6			1.3
Children's and young people's health	7 Smoking status at time of delivery	715	14.2	12.7	30.8			2.3
	8 Breastfeeding initiation	3,731	74.1	73.9	40.8			94.7
	9 Obese children (Year 6)	713	21.1	18.9	27.3			10.1
	10 Alcohol-specific hospital stays (under 18)	20	25.7	44.9	126.7		\rightarrow	11.9
	11 Under 18 conceptions	198	32.9	27.7	52.0			8.8
Adults' health and lifestyle	12 Smoking prevalence	n/a	20.5	19.5	30.1	<u> </u>		8.4
	13 Percentage of physically active adults	n/a	50.0	56.0	43.8			68.5
	14 Obese adults	n/a	19.6	23.0	35.2		0	11.2
	15 Excess weight in adults	470	57.0	63.8	75.9			45.9
Disease and poor health	16 Incidence of malignant melanoma	24	8.8	14.8	31.8			3.6
	17 Hospital stays for self-harm	488	137.4	188.0	596.0		0	50.4
	18 Hospital stays for alcohol related harm	1,988	717	637	1,121			365
	19 Drug misuse	2,704	12.1	8.6	26.3			8.0
	20 Recorded diabetes	24,551	8.4	6.0	8.7			3.5
	21 Incidence of TB	64	57.9	15.1	112.3	•	>	0.0
	22 Acute sexually transmitted infections	3,016	915	804	3,210		>	162
	23 Hip fractures in people aged 65 and over	218	513	568	828			403
Life expectancy and causes of death	24 Excess winter deaths (three year)	115	15.2	16.5	32.1			-3.0
	25 Life expectancy at birth (Male)	n/a	77.0	79.2	74.0			82.9
	26 Life expectancy at birth (Female)	n/a	81.8	83.0	79.5			86.6
	27 Infant mortality	34	6.4	4.1	7.5			0.7
	28 Smoking related deaths	366	293	292	480			172
	29 Suicide rate	32	10.3	8.5				
	30 Under 75 mortality rate: cardiovascular	208	108.5	81.1	144.7			37.4
	31 Under 75 mortality rate: cancer	293	151	146	213			106
	32 Killed and seriously injured on roads	90	27.3	40.5	116.3	♦		11.3

Indicator Notes

1 % people in this area living in 20% most deprived areas in England, 2010 2 % children (under 16) in families receiving means-tested benefits & low income, 2011 3 Crude rate per 1,000 households, 2012/13 4 % key stage 4, 2012/13 5 Recorded violence against the person crimes, crude rate per 1,000 population, 2012/13 6 Crude rate per 1,000 population aged 16-64, 2013 7 % of women who smoke at time of delivery, 2012/13 8 % of all mothers who breastfeed their babies in the first 48hrs after delivery, 2012/13 9 % school children in Year 6 (age 10-11), 2012/13 10 Persons under 18 admitted to hospital due to alcohol-specific conditions, crude rate per 100,000 population, 2010/11 to 2012/13 (pooled) 11 Under-18 conception rate per 1,000 females aged 15-17 (crude rate) 2012 12 % adults aged 18 and over, 2012 13 % adults achieving at least 150 mins physical activity per week, 2012 14 % adults classified as obese, Active People Survey 2012 15 % adults classified as overweight or obese, Active People Survey 2012 16 Directly age standardised rate per 100,000 population, aged under 75, 2009-2011 17 Directly age sex standardised rate per 100,000 population, 2012/13 18 The number of admissions involving an alcohol-related primary diagnosis or an alcohol-related external cause, directly age standardised rate per 100,000 population, 2012/13 19 Estimated users of opiate and/or crack cocaine aged 15-64, crude rate per 1,000 population, 2010/11 20 % people on GP registers with a recorded diagnosis of diabetes 2012/13 21 Crude rate per 100,000 population, 2010-2012 22 Crude rate per 100,000 population, 2012/13 24 Ratio of excess winter deaths (observed winter deaths minus expected deaths based on non-winter deaths) to average non-winter deaths 1.08.09-31.07.12 25 At birth, 2010-2012 26 At birth, 2010-2012 27 Rate per 1,000 live births, 2010-2012 28 Directly age standardised rate per 100,000 population aged under 75, 2010-2012 31 Directly age standardised rate per 100,000 population aged under 75, 2010-2012 32 Rate per 100,000 popul

More information is available at www.healthprofiles.info Please send any enquiries to healthprofiles@phe.gov.uk

© Crown copyright, 2014. You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence